

## Stroke

Simis, M. et al. Neurophysiologic predictors of motor function in stroke. Restor Neurol Neurosci. 2015; 34(1),45-54

Michelson, E.A. et al. Identification of acute stroke using quantified brain electrical activity. Acad Emerg Med. 2015 Jan;22(1), 67-72

Petti, M. et al. Individual cortical connectivity changes after stroke: a resampling approach to enable statistical assessment at single-subject level. Conf Proc IEEE Eng Med Biol Soc. 2014;2014,2785-2788

Schleiger, E. et al. Frontal EEG delta/alpha ratio and screening for post-stroke cognitive deficits: the power of four electrodes. Int J Psychophysiol. 2014 Oct;94(1),19-24

Song, Y. et al. Background rhythm frequency and theta power of quantitative EEG analysis: predictive biomarkers for cognitive impairment post-cerebral infarcts. Clin EEG Neurosci. 2015 Apr;46(2),142-146

Doruk, D. et al. Neurophysiologic Correlates of Post-stroke Mood and Emotional Control. Front Hum Neurosci. 2016 Aug 30;10,428

Alawieh, A., Zhao, J., Feng, W. Factors affecting post-stroke motor recovery: Implications on neurotherapy after brain injury. Behav Brain Res. 2016 Aug 13. 4328(16),30541-30551

Rozelle, G.R., Budzynski, T.H. Neurotherapy for stroke rehabilitation: a single case study. Biofeedback Self Regul.1995 Sep; 20(3),211-228

Schleiger, E. et al. Poststroke QEEG informs early prognostication of cognitive impairment. Psychophysiology. 2017 Feb;54(2),301-309

Finnigan, S., Wong, A., Read, S. Defining abnormal slow EEG activity in acute ischaemic stroke: Delta/alpha ratio as an optimal QEEG index. Clin Neurophysiol. 2016 Feb;127(2),1452-1459